

CONSTRUCTION OF DOUBLE COSET SYSTEM OF A COXETER GROUP AND ITS APPLICATIONS TO BRUHAT GRAPHS

MASATO KOBAYASHI*

ABSTRACT. We develop combinatorics of parabolic double cosets in finite Coxeter groups as a follow-up of recent articles by Billey-Konvalinka-Petersen-Slofstra-Tenner and Petersen. (1) We construct a double coset system as a generalization of a two-sided analogue of a Coxeter complex and present its order structure with its local dimension function on certain connected components. As applications of double cosets to Bruhat graphs, we also prove: (2) every parabolic double coset is regular, (3) invariance of degree on Bruhat graph on lower intervals as an analogy of the one for Kazhdan-Lusztig polynomials, (4) every noncritical Bruhat interval satisfies out-Eulerian property.

CONTENTS

1. Introduction	2
1.1. double cosets	2
1.2. main results	3
1.3. Organization	3
2. Preliminaries on Coxeter groups	3
2.1. weak, two-sided, Bruhat orders	3
2.2. Bruhat graphs	4
2.3. double cosets	4
2.4. presentations of a double coset	5
3. Double coset system	6
3.1. Petersen's two-sided analogue of the Coxeter complex	6
3.2. double coset system	7
3.3. variants of descent numbers	7
3.4. local structure: dimension, fiber, boolean complex	11
3.5. relation between Ξ and Δ	12
3.6. example	13
3.7. global structure: adjacent components	14

Date: July 27, 2019.

2010 Mathematics Subject Classification. Primary:20F55; Secondary:51F15.

Key words and phrases. Bruhat order, Bruhat graph, Coxeter group, Deodhar inequality, double cosets, Kazhdan-Lusztig polynomial, Poincaré polynomial.

*Department of Engineering, Kanagawa University, Japan.

3.8. theorem	14
4. Bruhat graphs on Bruhat intervals	15
4.1. Carrell-Peterson's result	15
4.2. regularity of Bruhat graphs	17
4.3. out-Eulerian property	21
5. Further remarks	22
5.1. four-variable Eulerian polynomials	22
5.2. left, right, central Poincaré polynomials	23
5.3. new enumeration problems on Bruhat graphs	24
References	25

1. INTRODUCTION

1.1. **double cosets.** Every Coxeter system possesses graded poset structures simultaneously as a (left/right) weak order and Bruhat order. These orders often show up in many topics: Coxeter complex, hyperplane arrangement, Eulerian polynomials, cluster algebras, Kazhdan-Lusztig polynomials and so on. When we consider two weak orders together (two-sided order), interesting interactions come into play which we must carefully analyze. One such example is to investigate structures of “double cosets” such as Solomon algebra or contingency tables. Recently, there are new developments in this topic:

- a two-sided analogue of a Coxeter complex and Eulerian polynomials: Petersen [13, 14] in 2018 and 2013.
- enumeration of double cosets with its minimal representative fixed: Billey-Konvalinka-Petersen-Slofstra-Tenner [2] in 2018.

Table 1. Three systems on W

one-sided Coxeter system	two-sided analogue	double coset system
$\Sigma(W)$	$\Xi(W)$	$\Delta(W)$
one-sided cosets	marked double cosets	double cosets
Tits	Hultman, Petersen	not studied
simplicial complex	boolean complex	?
$\dim(xW_I) = S \setminus I - 1$	$\dim(I, X, J) = S \setminus I + S \setminus J - 1$	?
$\max \dim = n - 1$	$\max \dim = 2n - 1$	?

1.2. main results. The aim of this article is to study combinatorics of parabolic double cosets and Bruhat graphs in finite Coxeter groups as a follow-up of their papers. We prove four main results as theorems:

- (1) Theorem 3.17: we construct a double coset system as a two-sided analogue of Coxeter complex and present several its order structures together with one-sided Coxeter complex and Petersen's analogue (Table 1).
- (2) Theorem 4.10: every parabolic double coset is regular,
- (3) Theorem 4.11: invariance of degree on Bruhat graph as an analogy of the one for Kazhdan-Lusztig polynomials,
- (4) Theorem 4.14: every noncritical Bruhat interval satisfies out-Eulerian property.

1.3. Organization. Section 2 gives basic ideas and facts on Coxeter systems. In Section 3, we construct a double coset system and show several results on its order structures. In Section 4, as applications of the idea double cosets (as Bruhat intervals), we prove three theorems on degree of a vertex on Bruhat graphs. In Section 5, we record some ideas and open problems for our research in the future.

2. PRELIMINARIES ON COXETER GROUPS

Throughout this article, we denote by $W = (W, S, T, \ell, \leq)$ a Coxeter system with W the underlying Coxeter group, S its Coxeter generators, T the set of its reflections, ℓ the length function, $\leq$ Bruhat order. Moreover, assume that W is finite of rank $n = |S|$. Unless otherwise noticed, symbols u, v, w, x, y are elements of W , $r, s \in S$, $t \in T$, e is the group-theoretic unit of W and I, J are subsets of S . The symbol $\ell(u, v)$ means $\ell(v) - \ell(u)$ for $u \leq v$, $x \triangleleft y$ a cover relation in a poset, and $w = u * v$ a reduced factorization of w : $w = uv$ as a group element, and $\ell(w) = \ell(u) + \ell(v)$.

2.1. weak, two-sided, Bruhat orders. We begin with basic definitions on partial orders on W .

Definition 2.1. Write

- (1) $u \triangleleft_L v$ if $\ell(u, v) = 1$ and $v = su$ for some $s \in S$.
- (2) $u \triangleleft_R v$ if $\ell(u, v) = 1$ and $v = us$ for some $s \in S$.
- (3) $u \triangleleft_{LR} v$ if $u \triangleleft_L v$ or $u \triangleleft_R v$.
- (4) $u \triangleleft v$ if $\ell(u, v) = 1$ and $v = tu$ for some $t \in T$ (equivalently, $v = ut'$ for some $t' \in T$).

(Further, we sometimes write $u \triangleleft_{2LR} v$ to mean $u \triangleleft_L v$ and $u \triangleleft_R v$.) Define four partial orders on W , *left weak order* ($\leq_L$), *right weak order* ($\leq_R$), *two-sided order* ($\leq_{LR}$) and *Bruhat order* ($\leq$), as transitive closure of those four binary relations on W . The interval notation is, for example,

$$[u, w]_{LR} = \{v \in W \mid u \leq_{LR} v \leq_{LR} w\}.$$

Say s is a *left (right) descent* of w if $\ell(sw) < \ell(w)$ ($\ell(ws) < \ell(w)$). Say s is a *left (right) ascent* of w if $\ell(sw) > \ell(w)$ ($\ell(ws) > \ell(w)$). We denote these sets of descents and ascents as follows:

$$\begin{aligned} D_L(w) &= \{s \in S \mid \ell(sw) < \ell(w)\}, \\ D_R(w) &= \{s \in S \mid \ell(ws) < \ell(w)\}, \\ A_L(w) &= \{s \in S \mid \ell(sw) > \ell(w)\} (= S \setminus D_L(w)), \\ A_R(w) &= \{s \in S \mid \ell(ws) > \ell(w)\} (= S \setminus D_R(w)). \end{aligned}$$

Also, the set of *left, right inversions* are

$$\begin{aligned} T_L(w) &= \{t \in T \mid \ell(tw) < \ell(w)\}, \\ T_R(w) &= \{t \in T \mid \ell(wt) < \ell(w)\}. \end{aligned}$$

2.2. Bruhat graphs.

Definition 2.2. The *Bruhat graph* of W is a directed graph for vertices $w \in W$ and for edges $u \rightarrow v$. For each subset $V \subseteq W$, we can also consider the induced subgraph with the vertex set V (Bruhat subgraph).

For convenience, we say an edge $x \rightarrow y$ is *short* if $\ell(x, y) = 1$ (i.e., $x \triangleleft y$). In other words, the short Bruhat graph for V is the directed version of the Hasse diagram of V .

Each subset of W can be regarded as a subposet under several kinds of graded partial orders (left weak, right weak, two-sided or Bruhat order).

2.3. double cosets. Most of our results rely on the following important property of Bruhat order:

Fact 2.3. Let $u < w$. If $s \in A_L(u) \cap D_L(w)$, then $su \leq w$ and $u \leq sw$ (Lifting Property). Consequently, if $v \in [u, w]$ and $s \in A_L(u) \cap D_L(w)$, then $sv \in [u, w]$.

The right version of this property also holds.

Definition 2.4. Let $I \subseteq S$. By W_I we mean the (standard) *parabolic subgroup* of W generated by I . A subset X in W is a *parabolic double coset* if

$$X = W_I x W_J = \{uxv \mid u \in W_I, v \in W_J\}$$

for some $x \in W$ and $I, J \subseteq S$.

In particular, every singleton set is a parabolic double coset itself while the empty set is not. By simply a coset or double coset, we mean a parabolic double coset hereafter.

Fact 2.5. Each double coset X has the representative of maximal and minimal length: there exists a unique pair $(x_0, x_1) \in X \times X$ such that

$$\ell(x_0) \leq \ell(x) \leq \ell(x_1)$$

for all $x \in X$.

Observation 2.6. A double coset X is (by construction) an interval in LR order. It is thus meaningful to write

$$X = "[x_0, x_1]_{LR}" = \{x \in W \mid x_0 \leq_{LR} x \leq_{LR} x_1\},$$

with $x_0 = \min X, x_1 = \max X$. Observe that if $X = W_I x W_J = [x_0, x_1]_{LR}$, then $I \subseteq A_L(x_0) \cap D_L(x_1)$ and $J \subseteq A_R(x_0) \cap D_R(x_1)$ otherwise x_0, x_1 cannot be the extremal elements of X . The *length* of a double coset X is $\ell(X) = \ell(\min X, \max X)$. Each $x \in X$ can be written as $x = u * x_0 * v$ for *some* $u \in W_I, v \in W_J, I, J \subseteq S$ with $I \subseteq A_L(x_0) \cap D_L(x_1)$ and $J \subseteq A_R(x_0) \cap D_R(x_1)$. Call u a left part of x , v a right part of x , and x_0 the central part of x (in X).

2.4. presentations of a double coset. We just introduced a double coset as a set in the form $X = W_I x W_J$. If $J = \emptyset$ at the extreme case, then $X = W_I x W_J = W_I x$ is an ordinary left coset. However, it is worth mentioning that the double coset $W_I x W_J$ may be equal to $W_I x$ (as sets) even if $J \neq \emptyset$ (or to $x W_J$ even if $I \neq \emptyset$). For example, the whole W is itself a double coset $W_S w_0 W_S$ and furthermore

$$W = W_S w_0 W_S = W_I e W_S = W_S x W_J.$$

Hence there are many ways to express a double coset with a certain choice of x and I, J .

Definition 2.7. Let X be a double coset. Say a triple (I, x, J) is a *presentation* of X if $X = W_I x W_J$.

Proposition 2.8. Say a presentation (I, x, J) of X is *maximal* if whenever

$$X = W_{I'} x' W_{J'},$$

then $I' \subseteq I, J' \subseteq J$.

Billey-Konvalinka-Petersen-Slofstra-Tenner [2, Proposition 3.7] proved that there is a unique maximal presentation for each double coset:

Fact 2.9. Let


$$x_0 = \min X, x_1 = \max X,$$

$$M_L(X) = A_L(x_0) \cap D_L(x_1),$$

$$M_R(X) = A_R(x_0) \cap D_R(x_1).$$

Then, $(M_L(X), x_1, M_R(X))$ is a unique maximal presentation of X .

Hence this is *the* maximal presentation of X . Similarly, we can talk about a minimal presentation of X in the following sense: a presentation (I, x, J) of X is *minimal* if whenever (I', x', J') is a presentation of X and $I' \subseteq I, J' \subseteq J$, then $I' = I, J' = J$. However, for a given X , there may exist more than one minimal presentation.

Figure 1. poset structures of $\Xi(A_1)$ and $\Delta(A_1)$ 

3. DOUBLE COSET SYSTEM

3.1. Petersen's two-sided analogue of the Coxeter complex. We first review Petersen's two-sided analogue of the Coxeter complex of W [13]. Motivated by Hultman [11], he constructed it as a collection of *marked double cosets*:

$$\Xi = \Xi(W) = \{(I, W_I x W_J, J) \mid x \in W, I, J \subseteq S\}.$$

He then introduced a partial order $(I, X, J) \leq_{\Xi} (I', X', J')$ by $I \supseteq I'$, $J \supseteq J'$ and $X \supseteq X'$. Each face (element) $F = (I, X, J)$ is colored by

$$\text{col}(F) = (S \setminus I, S \setminus J)$$


so that $\dim(F) = |S \setminus I| + |S \setminus J| - 1$.

Fact 3.1 ([13, Theorem 3]). For any Coxeter system (W, S) with $|S| = n < \infty$, we have the following.

- (1) The complex Ξ is a balanced boolean complex of dimension $2n - 1$.
- (2) The facets (maximal faces) of Ξ are in bijection with the elements of W , and the Coxeter complex Σ is a relative subcomplex of Ξ .
- (3) The complex Ξ is shellable and any linear extension of the two-sided weak order on W gives a shelling order for Ξ .
- (4) If W is finite then Ξ is contractible.
- (5) If W is infinite,
 - (a) the geometric realization of Ξ is a sphere, and
 - (b) a refined h -polynomial of Ξ is the *two-sided W -Eulerian polynomial*,

$$h(\Xi, s, t) = \sum_{w \in W} s^{\text{des}_L(w)} t^{\text{des}_R(w)}$$

where $\text{des}_L(w)$ denotes the number of left descents of w and $\text{des}_R(w)$ denotes the number of right descents of w .

Figure 2. the boolean complex $\Xi(A_2)$ (a copy of [13, Figure 1])

3.2. double coset system.

Definition 3.2. Define $\Delta = \Delta(W)$ be the set of all double cosets of W . Introduce a partial order on $\Delta(W)$ by the reverse of containment:


$$Y \leq_{\Delta} X \iff X \subseteq Y.$$

Unlike one-sided and two-sided Coxeter complexes, $(\Delta, \leq)$ is not necessarily a complex. However, it possesses some combinatorial structure with the “local dimension” function as we will see details below. For this reason, let us call the poset $(\Delta(W), \leq)$ the *double coset system* of W .

Example 3.3.

- (1) $W = A_1$ contains 5 marked cosets and 3 cosets (Figure 1).
- (2) $W = A_2$ contains 33 marked cosets and 19 cosets (Figures 2 and 3).

3.3. variants of descent numbers. Recall that $\dim(I, X, J) = |S \setminus I| + |S \setminus J| - 1$ the dimension function of Ξ . This is essentially counting a part of the ascent

Figure 3. double coset system $\Delta(A_2)$: 19 cosets

(descent) number of $\min x$ ($\max X$). How can we consider some analogue of this for Δ ? We will use the number of *weak coatoms* of a coset. For this purpose, let us prepare several definitions.

Table 2. variants of descent numbers on A_2

w	$d_{L1}(w)$	$d_{R1}(w)$	$d_{2LR}(w)$	$d(w)$	$\tilde{d}(w)$
123	0	0	0	0	0
132	0	0	1	1	2
213	0	0	1	1	2
231	1	1	0	2	2
312	1	1	0	2	2
321	0	0	2	2	4

Definition 3.4. A left descent $s \in D_L(w)$ is *small* if $w^{-1}sw \notin S$. Otherwise it is *large*. We use similar terminology for right descents.

More notation:

$$\begin{aligned}
 D_{L1}(w) &= \{s \in D_L(w) \mid w^{-1}sw \notin S\}, \\
 D_{L2}(w) &= \{s \in D_L(w) \mid w^{-1}sw \in S\}, \\
 D_{R1}(w) &= \{s \in D_R(w) \mid wsw^{-1} \notin S\},
 \end{aligned}$$

Table 3. two-sided and total descent numbers $d(w), \tilde{d}(w)$ over A_3

w	$d(w)$	$\tilde{d}(w)$	w	$d(w)$	$\tilde{d}(w)$	w	$d(w)$	$\tilde{d}(w)$	w	$d(w)$	$\tilde{d}(w)$
1234	0	0	2134	1	2	3124	2	2	4123	2	2
1243	1	2	2143	2	4	3142	3	3	4132	3	4
1324	1	2	2314	2	2	3214	2	4	4213	3	4
1342	2	2	2341	2	2	3241	3	4	4231	4	4
1423	2	2	2413	3	3	3412	2	2	4312	3	4
1432	2	4	2431	3	4	3421	3	4	4321	3	6

$$D_{R2}(w) = \{s \in D_R(w) \mid wsw^{-1} \in S\},$$

$$D_L(w) = D_{L1}(w) \cup D_{L2}(w), \quad D_R(w) = D_{R1}(w) \cup D_{R2}(w), \quad (\text{disjoint})$$

$$D_{2LR}(w) = \{(r, s) \in D_{L2}(w) \times D_{R2}(w) \mid rw = ws\}.$$

Observe that $|D_{L2}(w)| = |D_{R2}(w)| = |D_{2LR}(w)|$.

Left small descent number, left large descent number:

$$d_{L1}(w) = |D_{L1}(w)|, d_{L2}(w) = |D_{L2}(w)|,$$

Right small descent number, right large descent number:

$$d_{R1}(w) = |D_{R1}(w)|, d_{R2}(w) = |D_{R2}(w)|.$$

Two-sided descent number:

$$d(w) = d_{L1}(w) + d_{R1}(w) + d_{2LR}(w).$$

Total descent number:

$$\tilde{d}(w) = d_{L1}(w) + d_{R1}(w) + 2d_{2LR}(w) (= |D_L(w)| + |D_R(w)|).$$

Clearly, $d(w) \leq \tilde{d}(w) \leq 2d(w)$. Table 2 shows some examples.

Our method is to investigate double cosets with its maximal representative some fixed element (opposite to Billey et.al and Petersen). Let

$$\Delta(w) = \{X \in \Delta \mid \max X = w\}$$

and $\delta(w) = |\Delta(w)|$. Call $\Delta(w)$ the w -component of Δ . Naturally, $\Delta = \cup_{w \in W} \Delta(w)$ and the union is disjoint. Note $\min \Delta(w) = W_{D_L(w)} w W_{D_R(w)}$ and $\max \Delta(w) = \{w\}$. Although Billey et. al found the enumeration formula [2, Theorem 1.2] with the ‘‘marine model’’, we simply give some upper bound on $\delta(w)$ here.

Proposition 3.5.

$$\delta(w) \leq 2^{\tilde{d}(w)}.$$

Proof. Every coset X with $\max X = w$ has a marked coset expression $(I, W_I w W_J, J)$ with some $I \subseteq D_L(w), J \subseteq D_R(w)$. Notice that $D_L(w) = D_{L1}(w) \cup D_{L2}(w)$ and $D_R(w) = D_{R1}(w) \cup D_{R2}(w)$ are disjoint sums as introduced above. Thus these I, J can be expressed uniquely as

$$I = I_1 \cup I_2, \quad I_1 \subseteq D_{L1}(w), I_2 \subseteq D_{L2}(w),$$

$$J = J_1 \cup J_2, \quad J_1 \subseteq D_{R1}(w), J_2 \subseteq D_{R2}(w).$$

Hence

$$\delta(w) \leq |2^{D_{L1}(w)}| |2^{D_{L2}(w)}| |2^{D_{R1}(w)}| |2^{D_{R2}(w)}| = 2^{d_{L1}(w)} 2^{d_{L2}(w)} 2^{d_{R1}(w)} 2^{d_{R2}(w)} = 2^{\tilde{d}(w)}.$$

□

For example, let $w = 54312$. We see that

$$\begin{aligned} D_L(w) &= \{s_2, s_3, s_4\}, D_R(w) = \{s_1, s_2, s_3\}, \\ D_{L1}(w) &= \{s_2\}, D_{L2}(w) = \{s_3, s_4\}, D_{R2}(w) = \{s_1, s_2\}, D_{R1}(w) = \{s_3\}, \\ \tilde{d}(w) &= d_{L1}(w) + d_{L2}(w) + d_{R2}(w) + d_{R1}(w) = 1 + 2 + 2 + 1 = 6. \end{aligned}$$

Therefore, $\delta(w) \leq 2^6 = 32$.

Example 3.6. Figure 3 illustrates

$$\delta(123) = 1, \delta(213) = \delta(132) = 2, \delta(231) = \delta(312) = 4,$$

and in particular $\delta(321) = 6$ as boxed cosets show.

Example 3.7. $W = A_3$ has 167 cosets and 281 marked cosets [OEIS A260700, A120733]. Thanks to Table 3, we can check that

$$167 = \sum_{w \in A_3} \delta(w) \leq \sum_{w \in A_3} 2^{\tilde{d}(w)} \leq 1 \cdot 2^0 + 10 \cdot 2^2 + 2 \cdot 2^3 + 10 \cdot 2^4 + 1 \cdot 2^6 = 249 < 281.$$

3.4. local structure: dimension, fiber, boolean complex.

Definition 3.8. The set of weak coatoms for w is

$$C(w) = \{v \in W \mid v \triangleleft_{LR} w\}.$$

Similarly, the set of weak coatoms for a double coset $X = [x_0, x_1]_{LR}$ is

$$C(X) = \{v \in X \mid v \triangleleft_{LR} x_1\}.$$

Let $d(X) = |C(X)|$ (and $d(w) = |C(w)|$ as defined before). Moreover, let $\tilde{d}(X) = |M_L(X)| + |M_R(X)|$. Call $d(X)$ the *two-sided descent number* of X and $\tilde{d}(X)$ the *total descent number* of X .

Observation 3.9. For each $X \in \Delta(w)$, we have the following:

- (1) $0 \leq d(X) \leq n, 0 \leq \tilde{d}(X) \leq 2n$.
- (2) $d(X) = 0 \iff \tilde{d}(X) = 0$.
- (3) $d(X) = n \iff \tilde{d}(X) = 2n$.
- (4) $d(X) \leq \tilde{d}(X) \leq 2d(X)$.
- (5) $d(X) \leq d(w), \tilde{d}(X) \leq \tilde{d}(w)$.
- (6) If (I, X, J) is a presentation of X , then $d(X) \leq |I| + |J| \leq \tilde{d}(X)$.

Definition 3.10. The *local dimension* of $X \in \Delta(w)$ is

$$\dim_{\Delta(w)}(X) = d(w) - d(X) - 1.$$

If $X, Y \in \Delta(w)$ and $X \leq Y$, then $Y \subseteq X$ so that $C(Y) \subseteq C(X)$, $d(Y) \leq d(X)$ and $\dim_{\Delta(w)}(X) \leq \dim_{\Delta(w)}(Y)$. Thus, $\dim_{\Delta(w)}$ is a *weakly* increasing function on $(\Delta(w), \leq)$. At the extremal cases, we have $\dim_{\Delta(w)}(W_{D_L(w)}wW_{D_R(w)}) = d(w) - d(w) - 1 = -1$ is the minimum and $\dim_{\Delta(w)}(\{w\}) = d(w) - 0 - 1 = d(w) - 1$ is the maximum. In particular, call this $d(w) - 1$ the *dimension* of $\Delta(w)$. Moreover,

for any k with $-1 \leq k \leq d(w) - 1$, there exists some $X \in \Delta(w)$ such that $\dim_{\Delta(w)} X = k$ as easily shown.

3.5. relation between Ξ and Δ . Let $\Xi(w) := \{(I, X, J) \in \Xi \mid \max X = w\}$. This is a boolean subinterval in Ξ of rank $\tilde{d}(w)$ as seen from the proof of Proposition 3.5 (again, Petersen proved essentially the same result on marked cosets with minimal representative fixed [13, Theorem 9]); hence vertices $\Xi(w)$ and covering edges $\{(I, X, J) \triangleleft (I', X', J')\}$ form a connected subgraph in the Hasse diagram of Ξ . For this reason, we call $\Xi(w)$ the w -component of $\Xi(W)$. Clearly, $\Xi(W)$ is the disjoint union of these components:

$$\Xi(W) = \bigcup_{w \in W} \Xi(w).$$

Observation 3.11. The natural projection $\pi : \Xi \rightarrow \Delta$ by $\pi((I, X, J)) = X$ is weakly order-preserving.

$$(I, X, J) \leq (I', X', J') \implies \pi(I, X, J) \leq \pi(I', X', J').$$

Observe in particular that $\pi(\Xi(w)) = \Delta(w)$.

Definition 3.12. As an analogy of $\dim_{\Delta(w)} X = d(w) - d(X) - 1$, define the *local dimension function* on $\Xi(w)$ by

$$\dim_{\Xi(w)}(I, X, J) = \tilde{d}(w) - (|I| + |J|) - 1.$$

By definition of marked cosets, $\dim_{\Xi(w)}$ is a *strictly* increasing function on $(\Xi(w), \leq)$. At the extremal cases,

$$\dim_{\Xi(w)}(D_L(w), W_{D_L(w)}wW_{D_R(w)}, D_R(w)) = \tilde{d}(w) - (|D_L(w)| + |D_R(w)|) - 1 = -1$$

is the minimum and $\dim_{\Xi(w)}(\emptyset, \{w\}, \emptyset) = \tilde{d}(w) - 1$ is the maximum. This is indeed the dimension function of $\Xi(w)$ as a boolean complex.

The following proposition describes the relation of two local dimension functions via the projection:

Proposition 3.13. Let $(I, X, J) \triangleleft (I', X', J')$ in $\Xi(w)$ so that $\dim_{\Xi(w)}(I', X', J') - \dim_{\Xi(w)}(I, X, J) = 1$. Then, $\dim_{\Delta(w)} \pi(I', X', J') - \dim_{\Delta(w)} \pi(I, X, J) \in \{0, 1\}$. Consequently, if $(I, X, J) \leq (I', X', J')$ and $\dim_{\Xi(w)}(I', X', J') - \dim_{\Xi(w)}(I, X, J) = k$, then $\dim_{\Delta(w)} \pi(I', X', J') - \dim_{\Delta(w)} \pi(I, X, J) \in \{0, 1, 2, \dots, k\}$.

Proof. Let $(I, X, J) \triangleleft (I', X', J')$ in $\Xi(w)$. By definition, this means $X = W_I w W_J$, $X' = W_{I'} w W_{J'}$, $\max X = w = \max X'$ and either

- (1) $I = I' \cup \{s\}$, $J = J'$ for some $s \in D_L(w)$, $s \notin I'$ or
- (2) $I = I'$, $J = J' \cup \{s\}$ for some $s \in D_R(w)$, $s \notin J'$.

Say, for the moment, (1) holds. Considering the labels of edges between w and those coatoms, we have

$$\begin{aligned} C(X) &= \{rw \mid r \in I\} \cup \{wr \mid r \in J\}, \\ C(X') &= \{rw \mid r \in I'\} \cup \{wr \mid r \in J'\}, \end{aligned}$$

and $C(X') \subseteq C(X) = C(X') \cup \{sw\}$. Note that sw may or may not be in $C(X)$. Hence, $d(X) - d(X') \in \{0, 1\}$, that is,

$$\dim_{\Delta(w)} \pi(I', X', J') - \dim_{\Delta(w)} \pi(I, X, J) \in \{0, 1\}.$$

It is quite similar to show this in the case (2). The last part is shown by induction. $\square$

We can say more on relation between Ξ and Δ through the projection π . Recall that a finite poset P is a *boolean complex* (*simplicial poset*) if

- (1) $\widehat{0} \in P$ ($\widehat{0} \leq x$ for all $x \in P$),
- (2) each lower interval $[\widehat{0}, x]$ is a boolean poset.

Lemma 3.14. For each $X \in \Delta$, the fiber $\pi^{-1}(X)$ is a boolean complex.

Proof. Say $w = \max X$ so that $X \in \Delta(w)$. By definition,

$$\pi^{-1}(X) = \{(I, X, J) \in \Xi \mid I, J \subseteq S\}.$$

Note that, in this poset, $\widehat{0} = (M_L(X), X, M_R(X))$ is the unique minimal element. Furthermore, for each $(I, X, J) \in \pi^{-1}(X)$, the lower interval

$$[(M_L(X), X, M_R(X)), (I, X, J)]$$

is boolean since this is a subinterval of $\Xi(w)$ which is a boolean poset, and in fact every subinterval of a boolean poset is also boolean. $\square$

Proposition 3.15. Let $X \triangleleft \{w\}$ in Δ . Then X is covered by exactly two elements (and one of them is $\{w\}$).

Proof. Suppose $X \triangleleft \{w\}$ in Δ . Let (I, x, J) be the maximal presentation of X . Since X cannot be a singleton set ($X \triangleleft \{w\}$), we must have $|I| + |J| \geq 1$. Say $|I| \geq 1$ and choose $s \in I$. Then $X = W_I x W_J = W_I w W_J$ ($w \in X$) so that $\{w\} \subsetneq \{w, sw\} \subseteq X$ ($w \neq sw$). Since $\{w\}$ covers X , the set $\{w, sw\}$ must be X . In fact, $X = \{w, sw\} = W_{\{s\}} w$ is certainly a coset with $|X| = 2$. Hence X is covered by exactly two elements $\{w\}$ and $\{sw\}$. For the case $|J| \geq 1$, a similar argument is possible. $\square$

3.6. example. Let $W = A_2$, $S = \{s_1, s_2\}$ and $w = s_1 s_2 s_1$, the longest element. Figure 4 shows $\Xi(w)$ is a boolean interval of rank $4 = \widetilde{d}(w)$ with $2^{\widetilde{d}(w)} = 16$ marked cosets. It naturally splits into 6 boolean complexes of dimension 1, $-1, 0, 0, -1, -1$ (from the bottom), respectively:

$$|\Xi(321)| = 2^4 = 16 = 7 + 1 + 3 + 3 + 1 + 1.$$

These correspond to $6 = \delta(w)$ fibers for cosets in $\Delta(w)$. Observe also that $\pi : \Xi(w) \rightarrow \Delta(w)$ maps a boolean interval of rank 4 to a dihedral interval of rank 2 (six boxed cosets in Figure 3).

3.7. global structure: adjacent components. We discussed several “local” properties of Ξ and Δ . Here, we wish to present some “global” structures of such systems. First, we mention a less-known property on descent numbers.

Proposition 3.16. If $v \triangleleft_L w$, then $d_R(w) \in \{d_R(v), d_R(v) + 1\}$.

Proof. Suppose $v \triangleleft_L w$. By property of the left weak order, we have

$$T_R(v) \subsetneq T_R(w) = T_R(v) \uplus \{v^{-1}w\},$$

$$\underbrace{T_R(v) \cap S}_{D_R(v)} \subseteq \underbrace{T_R(w) \cap S}_{D_R(w)} = (T_R(v) \cap S) \uplus (\{v^{-1}w\} \cap S).$$

It follows that $D_R(w) = D_R(v) \uplus (\{v^{-1}w\} \cap S)$ and therefore $d_R(w) \in \{d_R(v), d_R(v) + 1\}$. $\square$

Now, consider the one-sided Coxeter complex $\Sigma = \Sigma(W) = \{xW_I \mid x \in W, I \subseteq S\}$. Let us call

$$\Sigma(w) = \{xW_I \mid x \in W, I \subseteq S, \max xW_I = w\}$$

the w -component of Σ . This is a boolean poset of rank $|D_R(w)| = d_R(w)$. Say components $\Sigma(v)$ and $\Sigma(w)$ are (left) *adjacent* if $v \triangleleft_L w$. Then, $d_R(w) \in \{d_R(v), d_R(v) + 1\}$ as shown above. Roughly speaking, adjacent components have close dimension.

We can do similar discussions for other systems. Say components $\Xi(v)$ and $\Xi(w)$ are (two-sided) *adjacent* if $v \triangleleft_{2LR} w$. By symmetry of left and right weak orders, $v \triangleleft_R w$ implies $d_L(w) \in \{d_L(v), d_L(v) + 1\}$. Consequently, if $v \triangleleft_{2LR} w$, then $d_L(w) = d_L(v) + 1, d_R(w) = d_R(v) + 1, d_{L1}(w) = d_{L1}(v), d_{L2}(w) = d_{L2}(v) + 1, d_{R1}(w) = d_{R1}(v), d_{R2}(w) = d_{R2}(v) + 1$ so that $\tilde{d}(w) = \tilde{d}(v) + 2$. In this way, adjacent components $\Xi(v), \Xi(w)$ have close dimension: $\tilde{d}(w) = \tilde{d}(v) + 2$.

Also, say components $\Delta(v)$ and $\Delta(w)$ are (two-sided) *adjacent* if $v \triangleleft_{2LR} w$. Then, for the same reason, $d(w) = d(v) + 1$.

It is not so obvious whether the whole Δ is ranked or not as Peterson pointed out [13, Remark 4]. We will study this point in future publication.

3.8. theorem. We summarize our results as a Theorem.

Theorem 3.17. The double coset system $(\Delta(W), \leq)$ has the following structures:

- (1) Set-theoretically, Δ is a disjoint union of $\Delta(w) = \{X \mid \max X = w\}$. In addition, each of them forms a connected subgraph of the Hasse diagram of $(\Delta, \leq)$; $\Delta(w)$ has the local dimension function $\dim_{\Delta(w)} : \Delta(w) \rightarrow$

- $\{-1, 0, 1, \dots, d(w) - 1\}$ which is weakly increasing and surjective. Moreover, $\Delta(w)$ has a unique minimal element $W_{D_L(w)}wW_{D_R(w)}$ of local dimension -1 . If $\Delta(v)$ and $\Delta(w)$ are adjacent, then $\dim \Delta(w) = \dim \Delta(v) + 1$.
- (2) Maximal elements of Δ are in bijection with elements of W .
 - (3) If a coset is covered by a maximal element in Δ , then it is covered by exactly two elements. (this is quite similar to the property Ξ is a pseudomanifold as Petersen proved)
 - (4) For each coset $X \in \Delta$, the fiber $\pi^{-1}(X)$ is a boolean complex. Moreover, for each $w \in W$,

$$\Xi(w) = \bigcup_{X \in \Delta(w)} \pi^{-1}(X)$$

gives a partition of a boolean interval of rank $\tilde{d}(w)$ into $\delta(w)$ boolean complexes.

It would be nice if we could apply some of these results (particularly (4)) to find another formula for $\delta(w)$.

4. BRUHAT GRAPHS ON BRUHAT INTERVALS

In this section, as applications of double cosets, we prove three theorems on degree of Bruhat graphs on Bruhat intervals. It is helpful for understanding our discussion to keep the following idea in mind: for $u, v \in [e, w]$, define $u \sim_w v$ if

$$W_{D_L(w)}uW_{D_R(w)} = W_{D_L(w)}vW_{D_R(w)}.$$

This gives a partition (an equivalent relation) of $[e, w]$:

$$[e, w] = \bigcup_{u \leq w} W_{D_L(w)}uW_{D_R(w)}.$$

4.1. Carrell-Peterson's result.

Definition 4.1. The *Poincaré polynomial* of w is

$$\mathcal{P}_w(q) = \sum_{v \leq w} q^{\ell(v)}.$$

The *average* of $\mathcal{P}_w(q)$ is $\mathcal{P}'_w(1)/\mathcal{P}_w(1)$.

Fact 4.2 ([5]). There exists a unique family of polynomials $\{P_{uw}(q) \mid u, w \in W\} \subseteq \mathbb{Z}[q]$ (*Kazhdan-Lusztig polynomials*) such that

- (1) $P_{uw}(q) = 0$ if $u \not\leq w$,
- (2) $P_{uw}(q) = 1$ if $u = w$,
- (3) $\deg P_{uw}(q) \leq (\ell(u, w) - 1)/2$ if $u < w$,

Figure 4. boolean interval $\Xi(s_1s_2s_1)$ of rank 4 and its partition into 6 boolean complexes as 6 fibers of $\pi : \Xi(s_1s_2s_1) \rightarrow \Delta(s_1s_2s_1)$. (I, J abbreviated and indication of cosets omitted)





Figure 5. an example of irregular Bruhat graphs



(4) if $u \leq w$, then

$$q^{\ell(u,w)} P_{uw}(q^{-1}) = \sum_{u \leq v \leq w} R_{uv}(q) P_{vw}(q),$$

where $\{R_{uv}(q) \mid u, v \in W\}$ are R -polynomials,

(5) $[q^0](P_{uw}) = 1$ if $u \leq w$.

Fact 4.3. Some invariance holds for a family of these polynomials: If $r \in D_L(w)$ and $s \in D_R(w)$, then $P_{ru,w}(q) = P_{uw}(q) = P_{us,w}(q)$. Notice that this statement includes even the $u \not\leq w$ case as $P_{ru,w}(q) = P_{uw}(q) = P_{us,w}(q) = 0$.

Fact 4.4 (Carrell-Peterson [7]). Suppose $P_{uw}(q)$ has nonnegative coefficients for all $u \leq w$. The following are equivalent:

- (1) $\mathcal{P}'_w(1)/\mathcal{P}_w(1) = \frac{1}{2}\ell(w)$.
- (2) Every $v \in [u, w]$ is incident to $\ell(u, w)$ edges.
- (3) $P_{uw}(q) = 1$ for all $u \leq w$.
- (4) $q^{\ell(w)} \mathcal{P}_w(q^{-1}) = \mathcal{P}_w(q)$.

Note that Carrell-Peterson's assumption is now true for all w in all Coxeter groups by Elias-Williamson [10].

4.2. regularity of Bruhat graphs. The statement (2) in Fact 4.4 is about degree and regularity of graphs. Let us see more details on this idea.

Definition 4.5. Let $\deg_V(v)$ denote the degree of v in Bruhat graph on the vertex set $V(\subseteq W)$. Define the *in-degree* and *out-degree* of v :

$$\text{in}_V(v) = \{u \in V \mid u \rightarrow v\},$$

$$\text{out}_V(v) = \{u \in V \mid v \rightarrow u\}.$$

For convenience, let $\text{in}_V(x) = \text{out}_V(x) = 0$ whenever $x \notin V$.

By definition, we have $\deg_V(v) = \text{in}_V(v) + \text{out}_V(v)$. For $V = W$, we simply write $\deg_W(v) = \deg(v)$, $\text{in}_W(v) = \text{in}(v) (= \ell(v))$ and $\text{out}_W(v) = \text{out}(v)$. For $V = [e, w]$, we also simply write $\deg_{[e,w]}(v) = \deg_w(v)$ and so on.

Question 4.6. What can we say about degree of Bruhat graph on a lower interval?

One remarkable fact is due to Deodhar [8], Dyer [9] and Polo [15]:

Fact 4.7 (Deodhar inequality [9]). We have

$$\deg_w(v) \geq \ell(w)$$

for all $v \in [e, w]$.

(Similar statement holds for general Bruhat intervals)

Now recall that a finite directed graph is *d-regular* if $\deg(v) = d$ for all vertex $v \in V$. It is *regular* if it is *d-regular* for some nonnegative integer d . For example, the Hasse diagram of a finite boolean poset is regular; the Bruhat graph of the whole $W = [e, w_0]$ is $|T|$ -regular.

The following fact shows some special characteristic of Bruhat graphs.

Fact 4.8. The following are equivalent:

- (1) $[e, w]$ is regular.
- (2) $[e, w]$ is $\ell(w)$ -regular.

Remark: at a glance, it may be possible that even if there exists some $v \in [e, w]$ such that $\deg(v) > \ell(w)$, the Bruhat graph $[e, w]$ still can be regular (i.e., $\deg(v)$ -regular), but that is not true; notice that degree of the top element w is exactly $\ell(w)$.

One-sided cosets $W_I x, xW_J$ are always regular ($\ell(w_0(I))$ -regular, $\ell(w_0(J))$ -regular where $w_0(I), w_0(J)$ are the longest elements of W_I, W_J). How about as short graphs? Indeed, $W_I x, xW_J$ are *short-regular* meaning its short Bruhat graph is regular ($|I|$ -regular, $|J|$ -regular to be precise). However, this does not hold for double cosets. A counterexample appears even in $W = A_2$ ($s_1 = (12), s_2 = (23)$):

$$X = W = W_{\{s_1, s_2\}} e W_{\{s_1, s_2\}}$$

is itself a coset. The short degree of $e = 123$ in X is 2 while the one of $v = s_1 = 213$ is 3.

Question 4.9 (Table 4). Is every double coset regular as a Bruhat graph?

The answer is indeed yes as shown below; this result might be proved in geometric method (such as theory of Schubert varieties [3] and Richardson varieties [4]), here we give a combinatorial proof.

Table 4. regularity of finite cosets

	$W_I x$	$x W_J$	$W_I x W_J$
as a short graph	regular	regular	not necessarily regular
as a Bruhat graph	regular	regular	?

Theorem 4.10. Every double coset X is regular. To be more precise, X is $\ell(X)$ -regular.

Proof. Let $X = [x_0, x_1]$ be a double coset. For $x \in X$, define

$$\begin{aligned} \text{in}(x) &= |\{w \in W \mid w \rightarrow x, w \notin X\}|, \\ \text{self}_\uparrow(x) &= |\{w \in W \mid w \rightarrow x, w \in X\}|, \\ \text{self}^\uparrow(x) &= |\{w \in W \mid x \rightarrow w, w \in X\}|. \end{aligned}$$

Observe that if $x = u * x_0 * v$ with $X = W_I x W_J$, $u \in W_I, v \in W_J$, then

$$\text{in}(x) = \ell(x_0), \quad \text{self}_\uparrow(x) = \ell(u) + \ell(v).$$

(to see this, consider a reduced factorization

$$x = a_1 \cdots a_l * b_1 \cdots b_m * c_1 \cdots c_n$$

for x with $a_i, b_j, c_k \in S$, $u = a_1 \cdots a_l, x_0 = b_1 \cdots b_m, v = c_1 \cdots c_n$ all reduced, $t_i = a_1 \cdots a_{i-1} a_i a_{i-1} \cdots a_1 \in T_L(u) \subseteq T_L(x)$ and $t'_i = c_n \cdots c_{n-i+2} c_{n-i+1} c_{n-i+2} \cdots c_n \in T_R(v) \subseteq T_R(x)$. It is easy to see $t_i x \neq x t'_i$ for all i .) It follows that

$$\ell(x) = \text{in}(x) + \text{self}_\uparrow(x),$$

$$\deg_X(x) = \text{self}_\uparrow(x) + \text{self}^\uparrow(x).$$

Now we need to show that for all $x, y \in X$, $\deg_X(x) = \deg_X(y)$. Since X is a graded poset, It is enough to show this for $(x, y) \in X^2$ with $x \triangleleft_{LR} y$. Say, for the moment, $x \triangleleft_L y, y = sx, s \in I \subseteq A_L(x_0) \cap D_L(x_1)$. Note that

$$\text{in}(x) = \ell(x_0) = \text{in}(y) \text{ and } \ell(y) = \ell(x) + 1.$$

We claim that for each $t \in T \setminus \{x^{-1}sx\}$, we have

$$x \rightarrow xt \in X \iff y \rightarrow yt \in X.$$

If $x \rightarrow xt \in X$ (in particular $t \notin T_L(x)$), then $y \rightarrow yt$ since $T_L(x) \subseteq T_L(y) = T_L(x) \cup \{x^{-1}sx\}$. Thus $t \notin T_L(y)$. Moreover,

$$yt = (sx)t = \underbrace{s}_{\in A_L(x_0) \cap D_L(x_1)} \underbrace{(xt)}_{\in X} \in X$$

by Lifting Property (and vice versa). We proved the claim. If $x \triangleleft_R y, y = xs, s \in J \subseteq A_R(x_0) \cap D_R(x_1)$, then the similar proof goes with t on the left side. Consequently,

$$\begin{aligned} \deg_X(y) - \deg_X(x) &= (\text{self}_\uparrow(y) + \text{self}^\uparrow(y)) - (\text{self}_\uparrow(x) + \text{self}^\uparrow(x)) \\ &= (\text{self}_\uparrow(y) - \text{self}_\uparrow(x)) + (\text{self}^\uparrow(y) - \text{self}^\uparrow(x)) \\ &= (\ell(y) - \text{in}(y)) - (\ell(x) - \text{in}(x)) - 1 \\ &= (\ell(y) - \ell(x)) - (\text{in}(y) - \text{in}(x)) \\ &= 0. \end{aligned}$$

Hence $\deg(x) = \deg(x_1) = \ell(X)$. □

The following is an analogy of invariance of Kazhdan-Lusztig polynomials: $P_{rv,w}(q) = P_{vw}(q) = P_{vs,w}(q)$ for all $v \in W, r \in D_L(w)$ and $s \in D_R(w)$.

Theorem 4.11. We have

$$\deg_w(rv) = \deg_w(v) = \deg_w(vs)$$

for all $v \in W, r \in D_L(w)$ and $s \in D_R(w)$.

Proof. It is enough to show that $\deg_w(v) = \deg_w(rv)$ for $v \in [e, w]$ and $r \in D_L(w)$. Replacing v by rv if necessary, we may assume that $v \rightarrow rv$. Split $\deg_w(v)$ as:

$$\deg_w(v) = \text{in}_w(v) + \text{out}_w(v).$$

It follows that

$$\text{in}_w(v) = \ell(v), \quad \text{in}_w(rv) = \ell(rv) = \ell(v) + 1.$$

Now we claim that for $t \in T \setminus \{v^{-1}rv\}$, we have

$$v \rightarrow vt \leq w \iff rv \rightarrow (rv)t \leq w.$$

If $v \rightarrow vt \leq w$ (in particular $t \notin T_R(v)$), then $rv \rightarrow rvt$ since

$$T_R(v) \subsetneq T_R(rv) = T_R(v) \cup \{v^{-1}rv\}.$$

Moreover, $rvt = r \underbrace{(vt)}_{\leq w} \leq w$ by Lifting Property. We showed ($\implies$) (and vice versa). We proved the claim. In addition, v is incident to exactly one more outgoing edge $v \rightarrow rv (= v(v^{-1}rv)) \leq w$. Hence

$$\text{out}_w(v) = \text{out}_w(rv) + 1.$$

Altogether, conclude that

$$\deg_w(v) = \text{in}_w(v) + \text{out}_w(v) = \ell(v) + \text{out}_w(rv) + 1 = \deg_w(rv).$$

□

4.3. out-Eulerian property. Recall the basic fact that every Bruhat interval is Eulerian. In particular,

$$\sum_{v \in [e, w]} (-1)^{\ell(v)} = 0$$

for $w \neq e$. For a proof of this, choose $t \in T \cap W_{D_L(w)}$. We see that $v \leftrightarrow tv$ is a perfect matching on $[e, w]$ as a consequence of Lifting Property. Moreover, as is well-known, $\ell(v, tv)$ is always odd. Hence

$$\sum_{v \in [e, w]} (-1)^{\ell(v)} = \sum_{v \leftrightarrow tv} ((-1)^{\ell(v)} + (-1)^{\ell(tv)}) = 0.$$

(Usually, we take t to be a simple reflection. However, it is not necessary.) In terms of Bruhat graphs, we can understand this Eulerian property as

$$\sum_{v \leq w} (-1)^{\text{in}_w(v)} = 0.$$

It is natural to wonder if the similar statement on out-degree holds. The point is: if $u \rightarrow v$ in $[e, w]$, is always $\text{out}_w(u) - \text{out}_w(v)$ odd? The answer is no; but this is far from obvious and not so often this idea has been mentioned in the literature.

Definition 4.12. Say an edge $u \rightarrow v$ in $[e, w]$ is *out-odd* if $\text{out}_w(u) - \text{out}_w(v)$ is odd. It is *out-even* if $\text{out}_w(u) - \text{out}_w(v)$ is even.

We can easily find an example of both kinds (Figure 5): $3124 \rightarrow 3214$ is out-odd in $[1324, 3412]$ since

$$\text{out}_{3412}(3124) - \text{out}_{3412}(3214) = 2 - 1 = 1$$

while $1324 \rightarrow 3124$ is out-even since

$$\text{out}_{3412}(1324) - \text{out}_{3412}(3124) = 4 - 2 = 2.$$

We will show that “out-Eulerian property” holds for some special class of Bruhat intervals:

Definition 4.13. We say that $(u, w) \in W \times W$ is a *critical pair* if

$$u \leq w, \quad D_L(w) \subseteq D_L(u) \text{ and } D_R(w) \subseteq D_R(u).$$

An interval $[u, w]$ is *critical* if (u, w) is a critical pair.

Now we have a simple classification:

$$\text{Bruhat intervals} \begin{cases} \text{critical} \\ \text{noncritical} \end{cases}$$

A trivial interval is critical; a lower interval $[e, w]$ ($w \neq e$) is noncritical; a double coset of length ≥ 1 is noncritical. Observe that if $[e, w]$ is noncritical, then there exists some $s \in S$ such that $s \in A_L(u) \cap D_L(w)$ or $s \in A_R(u) \cap D_R(w)$.

Theorem 4.14 (out-Eulerian Property). For every noncritical interval $[u, w]$, we have

$$\sum_{v \in [u, w]} (-1)^{\text{out}_{[u, w]}(v)} = 0.$$

Notice that $\text{out}_{[u, w]}(v) = \text{out}_w(v)$ for $v \in [u, w]$.

Proof. Consider the partition of $[e, w]$:

$$[e, w] = \bigcup_{v \in [e, w]} W_{D_L(w)} v W_{D_R(w)}.$$

Assume $[u, w]$ is noncritical. Then, (say left) $A_L(u) \cap D_L(w) \neq \emptyset$ and choose $t \in T \cap W_{D_L(w)}$. For each $v \in [u, w]$, $v \leftrightarrow tv$ is a perfect matching on $[u, w]$ as a consequence of Lifting Property, again. Moreover, $v \sim_w tv$ implies

$$\deg_w(v) = \deg_w(tv)$$

as proved in Theorem 4.11. It follows that

$$\text{in}_w(v) + \text{out}_w(v) = \text{in}_w(tv) + \text{out}_w(tv),$$

$$\ell(v) + \text{out}_w(v) = \ell(tv) + \text{out}_w(tv),$$

$$\ell(v) - \ell(tv) = \text{out}_w(tv) - \text{out}_w(v).$$

Since $\ell(v) - \ell(tv)$ is odd, so is $\text{out}_w(tv) - \text{out}_w(v)$. Conclude that

$$\begin{aligned} \sum_{v \in [u, w]} (-1)^{\text{out}_{[u, w]}(v)} &= \sum_{v \in [u, w]} (-1)^{\text{out}_w(v)} \\ &= \sum_{v \leftrightarrow tv} ((-1)^{\text{out}_w(v)} + (-1)^{\text{out}_w(tv)}) \\ &= 0. \end{aligned}$$

□

5. FURTHER REMARKS

We end with recording some ideas for subsequent research.

5.1. four-variable Eulerian polynomials. It is possible to consider the following W -Eulerian polynomial in four variables

$$A_W(t_1, t_2, t_3, t_4) = \sum_{w \in W} t_1^{d_{L1}(w)} t_2^{d_{L2}(w)} t_3^{d_{R2}(w)} t_4^{d_{R1}(w)}.$$

Notice that $t_1 = t_2 = t$ and $t_3 = t_4 = 1$ recovers classical W -Eulerian polynomial (Brenti [6]) and $t_1 = t_2 = s, t_3 = t_4 = t$ recovers two-sided W -Eulerian polynomial (Petersen [14]). What if $t_1 = t, t_2 = t_3 = t^{1/2}, t_4 = t$ (the generating function of *two-sided descent number*) or $t_1 = t_2 = t_3 = t_4 = t$ (the generating function of *total descent number*)?

This polynomial must satisfy $|\Delta(W)| \leq A_W(2, 2, 2, 2)$ and $A_W(t_1, t_2, t_3, t_4) = A_W(t_4, t_3, t_2, t_1)$ since $d_{L1}(w^{-1}) = d_{R1}(w)$, $d_{L2}(w^{-1}) = d_{R2}(w)$ and so on.

5.2. left, right, central Poincaré polynomials. As is well-known, Poincaré polynomials have nice factorization property with respect to cosets and quotients [5]. Here let us consider more variants of Poincaré polynomials. Recall that $w_0(I)$ means the longest element of W_I .

Definition 5.1. Define maps (*left, coleft, right, coright projections*) $L, \tilde{L}, R, \tilde{R} : W \rightarrow W$ by

$$\begin{aligned} L(x) &= w_0(D_L(x)), & \tilde{L}(x) &= L(x)^{-1}x, \\ R(x) &= w_0(D_R(x)), & \tilde{R}(x) &= xR(x)^{-1}. \end{aligned}$$

so that we have

$$x = L(x)\tilde{L}(x) \text{ and } \ell(x) = \ell(L(x)) + \ell(\tilde{L}(x))$$

$$x = \tilde{R}(x)R(x) \text{ and } \ell(x) = \ell(\tilde{R}(x)) + \ell(R(x)).$$

Call $L(w)$ ($\tilde{L}(w)$, $R(w)$, $\tilde{R}(w)$) the *left* (*coleft*, *right*, *coright*) *part* of w , and left length, left colength right length, right colength

$$\begin{aligned} \ell_L(x) &= \ell(L(x)), & \ell_{\tilde{L}}(x) &= \ell(\tilde{L}(x)), \\ \ell_R(x) &= \ell(R(x)), & \ell_{\tilde{R}}(x) &= \ell(\tilde{R}(x)). \end{aligned}$$

Definition 5.2. Let $C(w) = \min W_{D_L(w)}wW_{D_R(w)}$ be the *central projection*. $\ell_C(w) = \ell(C(w))$, $\ell_{\text{side}}(w) = \ell(w) - \ell_C(w)$: *central length* and *side length* of w .

For example, $w = 45312$ has a reduced word $s_2s_3s_2s_1s_4s_2s_3s_2$ with $D_L(w) = D_R(w) = \{s_2, s_3, s_2\}$. Thus,

$$45312 = \underbrace{s_2s_3s_2}_{L(w)} \underbrace{s_1s_4s_2s_3s_2}_{\tilde{L}(w)} = \underbrace{s_2s_3s_2s_1s_4}_{\tilde{R}(w)} \underbrace{s_2s_3s_2}_{R(w)} = s_2s_3s_2 \underbrace{s_1s_4}_{C(w)} s_2s_3s_2$$

$$\ell_L(w) = \ell_R(w) = 3, \ell_{\tilde{L}}(w) = \ell_{\tilde{R}}(w) = 5, \ell_C(w) = 2, \ell_{\text{side}}(w) = 6.$$

Left, Right, Central Poincare polynomials of w :

$$\begin{aligned} \mathcal{P}_w^L(q_1, q_2) &= \sum_{v \leq w} q_1^{\ell_L(v)} q_2^{\ell_{\tilde{L}}(v)}, \\ \mathcal{P}_w^R(q_1, q_2) &= \sum_{v \leq w} q_1^{\ell_{\tilde{R}}(v)} q_2^{\ell_R(v)}, \\ \mathcal{P}_w^C(q_1, q_2) &= \sum_{v \leq w} q_1^{\ell_C(v)} q_2^{\ell_{\text{side}}(v)}. \end{aligned}$$

In particular, $\mathcal{P}_w^L(q, q) = \mathcal{P}_w^R(q, q) = \mathcal{P}_w^C(q, q) = \mathcal{P}_w(q)$. Find these polynomials.

5.3. new enumeration problems on Bruhat graphs.

- (1) Let $V(w)$ be the vertex set of $[e, w]$ and $E(w) = \{u \rightarrow v \mid u, v \in [e, w]\}$ the set of all edges. Say a vertex v is *irregular* if $\deg_w(v) > \ell(w)$. Say an edge $u \rightarrow v$ is *irregular* if it is incident to an irregular vertex; see [12, Theorem 8.2] some relation between irregularity and edges of Bruhat graphs. The following rational numbers seem to be quite natural to “measure irregularity” of $[e, w]$:

$$\frac{|V_{\text{irr}}(w)|}{|V(w)|}, \frac{|E_{\text{irr}}(w)|}{|E(w)|},$$

However, these have not been studied. Compute some examples. Can these numbers be any rational number between 0 and 1?

- (2) When is an edge $u \rightarrow v$ in $[e, w]$ out-even or when not? Try Type A. Describe it in terms of reduced words, monotone triangles and pattern avoidance.
- (3) Further, the *in-out-Poincaré polynomial* of w is

$$\mathcal{P}_w^{\text{in-out}}(q_1, q_2) = \sum_{v \leq w} q_1^{\text{in}_w(v)} q_2^{\text{out}_w(v)}.$$

In particular, *out-Poincaré polynomial* of w is

$$\mathcal{P}_w^{\text{out}}(q) = \mathcal{P}_w^{\text{in-out}}(1, q) = \sum_{v \leq w} q^{\text{out}_w(v)}$$

as we showed that $\mathcal{P}_w^{\text{out}}(-1) = 0$ for $w \neq e$. Study these polynomials. When are they palindromic?

- (4) The following are equivalent [3]:
- (a) $[e, w]$ is irregular.
 - (b) There exists some $v \in [e, w]$ such that $\deg(v) > \ell(w)$.
 - (c) $\deg_w(e) > \ell(w)$.

The degree function in (c) is interesting:

$$\deg_w(e) = |\{v \in [e, w] \mid e \rightarrow v\}| = |\{t \in T \mid t \leq w \text{ (subword)}\}|.$$

Let $\lambda(w) = \deg_w(e)$. By definition, λ is *weakly* increasing in Bruhat order:

$$x \leq y \implies \lambda(x) \leq \lambda(y).$$

Observe also that $\lambda(e) = 0 = \ell(e)$, $\lambda(w_0) = |T| = \ell(w_0)$ and moreover $\lambda(w) \geq \ell(w)$ due to Deodhar inequality. Let us say w is *combinatorially smooth* [1] if $\lambda(w) = \ell(w)$. It is not so easy to predict when $\lambda(w) > \ell(w)$ as the example shows below:

$$\begin{array}{cccccccccccc} w & 1234 & \rightarrow & 1324 & \rightarrow & 3124 & \rightarrow & 3142 & \rightarrow & 3412 & \rightarrow & 4312 & \rightarrow & 4321 \\ \lambda(w) & 0 & \rightarrow & 1 & \rightarrow & 2 & \rightarrow & 3 & \rightarrow & 5 & \rightarrow & 5 & \rightarrow & 6 \end{array}$$

$$\lambda(3142) = |\{t \in T \mid t \leq s_2 s_1 s_3\}| = |\{s_1, s_2, s_3\}| = 3,$$

$$\begin{aligned} \lambda(3412) &= |\{t \in T \mid t \leq s_2 s_1 s_3 s_2\}| \\ &= |\{s_1, s_2, s_3, s_2 s_1 s_2, s_2 s_3 s_2\}| = 5, \end{aligned}$$

$$\begin{aligned} \lambda(4312) &= |\{t \in T \mid t \leq s_2 s_1 s_3 s_2 s_1\}| \\ &= |\{s_1, s_2, s_3, s_2 s_1 s_2, s_2 s_3 s_2\}| = 5, \end{aligned}$$

Discuss edges $v \rightarrow w$ such that $\lambda(v, w) = 0, 1$ or ≥ 2 .

REFERENCES

- [1] Abe-Billey, Consequences of the Lakshmibai-Sandhya Theorem: the ubiquity of permutation patterns in Schubert calculus and related geometry, Schubert calculus–Osaka 2012, 1-52, Adv. Stud. Pure Math., 71, Math. Soc. Japan, 2016.
- [2] Billey-Konvalinka-Petersen-Slofstra-Tenner, Parabolic double cosets in Coxeter groups, Electr. J. Combin. Volume 25 (2018), no.1, Paper 1.23, 66pp.
- [3] Billey-Lakshmibai, Singular loci of Schubert varieties, Progress in Mathematics, 182. Birkhäuser Boston, Inc., Boston, MA, 2000.
- [4] Billey-Coskun, Singularities of generalized Richardson varieties, Comm. Algebra 40 (2012), no. 4, 1466-1495.
- [5] Björner-Brenti, Combinatorics of Coxeter groups, Graduate Texts in Mathematics, vol. 231, Springer-Verlag, New York, 2005.
- [6] Brenti, q -Eulerian polynomials arising from Coxeter groups, Europ. J. Combin. 15 (1994), no. 5, 417-441.
- [7] Carrell, The Bruhat graph of a Coxeter group, a conjecture of Deodhar, and rational smoothness of Schubert varieties, Proc. Symp. Pure Math., Vol. 56, 1994, 53-61.
- [8] Deodhar, Local Poincaré duality and nonsingularity of Schubert varieties, Comm. Algebra 13 (1985), no. 6, 1379-1388.
- [9] Dyer, The nil Hecke ring and Deodhar's conjecture on Bruhat intervals, Invent. Math. 111 (1993), no.3, 571-574.
- [10] Elias-Williamson, The Hodge theory of Soergel bimodules, Ann. of Math. (2) 180 (2014), no. 3, 1089-1136.
- [11] Hultman, The combinatorics of twisted involutions in Coxeter groups, Trans. Amer. Math. Soc. 359 (2007), no. 6, 2787-2798.
- [12] Kobayashi, Inequalities on Bruhat graphs, R - and Kazhdan-Lusztig polynomials, J. Combin. Theory Ser. A 120 (2013), no.2, 470-482.
- [13] Petersen, a two-sided analogue of the Coxeter complex, Electr. J. Combin. 25 (2018), no. 4, Paper 4.64, 28pp.
- [14] Petersen, Two-sided Eulerian numbers via balls in boxes, Math. Mag. 86 (2013), no. 3, 159-176.
- [15] Polo, On Zariski tangent spaces of Schubert varieties, and a proof of a conjecture of Deodhar, Indag. Math. (N.S.) 5 (1994), no. 4, 483-493.

DEPARTMENT OF ENGINEERING, KANAGAWA UNIVERSITY, 3-27-1 ROKKAKU-BASHI, YOKOHAMA 221-8686, JAPAN.

E-mail address: masato210@gmail.com